

Algebra & Number Theory

Volume 16
2022
No. 5

Entirety of certain cuspidal Eisenstein series on Kac–Moody groups

Lisa Carbone, Kyu-Hwan Lee and Dongwen Liu



Entirety of certain cuspidal Eisenstein series on Kac–Moody groups

Lisa Carbone, Kyu-Hwan Lee and Dongwen Liu

Let G be an infinite-dimensional representation-theoretic Kac–Moody group associated to a nonsingular symmetrizable generalized Cartan matrix. We consider Eisenstein series on G induced from unramified cusp forms on finite-dimensional Levi subgroups of maximal parabolic subgroups. Under a natural condition on maximal parabolic subgroups, we prove the cuspidal Eisenstein series are entire on the full complex plane.

1. Introduction

Meromorphic continuation of Eisenstein series on reductive groups, established in the seminal work of Langlands [1976], has been a foundational basis for other developments in Langlands’ program. Most notably, the Langlands–Shahidi method (e.g., [Kim and Shahidi 2002; Kim 2003]) in the study of L -functions originates from the idea of exploiting analytic properties of Eisenstein series together with the fact that L -functions appear in the Fourier coefficients of Eisenstein series.

The extension of the theory of Eisenstein series to Kac–Moody groups has been developed with anticipation of its potential roles in some of the central problems in number theory [Bump et al. 1996; Shahidi 2005]. Indeed, starting with Garland’s pioneering work [1999; 2004; 2006], there has been a significant body of work [Garland 2007a; 2007b; 2007c; 2007d; 2017; 2011; Liu 2015] on Eisenstein series on affine Kac–Moody groups, including the function field case [Braverman and Kazhdan 2013; Kapranov 2000; Patnaik 2008; Lee and Lombardo 2014]. Beyond affine Kac–Moody groups, Eisenstein series on rank 2 symmetric hyperbolic Kac–Moody groups were studied in [Carbone et al. 2017]. For a more general class of Kac–Moody groups, in [Carbone et al. 2020], the absolute convergence of Eisenstein series is established for spectral parameters in the Godement range.

In the process of these developments, a striking difference between the finite-dimensional case and the affine case was observed in [Braverman and Kazhdan 2013; Garland et al. 2017], where the affine Eisenstein series induced from cusp forms on finite-dimensional Levi subgroups were shown to be *entire* and not just meromorphic as they are in the finite-dimensional case. This phenomenon is not restricted to the affine case, and it was shown in [Carbone et al. 2017] that cuspidal Eisenstein series on rank 2 symmetric hyperbolic Kac–Moody groups are also entire.

Carbone was supported in part by the Simons Foundation, Mathematics and Physical Sciences Collaboration Grants for Mathematicians, Award Number: 422182. Lee was partially supported by a grant from the Simons Foundation (#712100). Liu was partially supported by the Fundamental Research Funds for the Central Universities 2016QNA2002.

MSC2020: 11F70, 20G44.

Keywords: Eisenstein series, Kac–Moody groups, entirety.

To understand the analytic properties of cuspidal Eisenstein series on Kac–Moody groups, one may naturally ask: *For which parabolic subgroups are cuspidal Eisenstein series on Kac–Moody groups entire?* For a general maximal parabolic subgroup, even in the finite-dimensional case cuspidal Eisenstein series will not be entire, as is readily seen for the real analytic Eisenstein series on SL_n .

In this paper, we give a natural condition on parabolic subgroups (property RD, below and Section 4) and show that cuspidal Eisenstein series attached to a parabolic subgroup satisfying property RD is holomorphic on the full complex plane.

More precisely, let G be a representation-theoretic Kac–Moody group, and let \mathfrak{g} be the corresponding real Kac–Moody algebra with a fixed Cartan subalgebra \mathfrak{h} , Weyl group W and the set $\{\alpha_i\}_{i \in I}$ of simple roots, where I is the set of indices for the simple roots. We assume that \mathfrak{g} is infinite-dimensional and nonaffine since the other cases have been studied well as mentioned above.

The real group $G_{\mathbb{R}}$ has the Iwasawa decomposition $G_{\mathbb{R}} = UA^+K$ [De Medts et al. 2009], where U is a maximal pronipotent subgroup, A^+ is the connected component of a maximal torus, and K is a subgroup of $G_{\mathbb{R}}$ analogous to the maximal compact subgroup in the finite-dimensional theory. Denote by Γ the arithmetic subgroup of $G_{\mathbb{R}}$ (Section 2A), and by $\mathcal{C} \subset \mathfrak{h}$ the open fundamental chamber. Using the exponential map $\exp : \mathfrak{h} \rightarrow A^+$, set $A_{\mathcal{C}} = \exp \mathcal{C}$.

Let P be a maximal parabolic subgroup of $G_{\mathbb{R}}$ with fixed Levi decomposition and associated finite-dimensional Levi subgroup M , which is associated with a subset $\theta \subseteq I$. We denote by α_P the simple root associated to the one element index in $I \setminus \theta$. Let L be the derived subgroup of M . For a cusp form f on $(L \cap \Gamma) \backslash L$, we recall that f is unramified if f is right-invariant under the action of $L \cap K$.

For such an f , we define the Eisenstein series $E_f(s, g)$, $s \in \mathbb{C}$, $g \in G_{\mathbb{R}}$, in analogy with the classical case (see (3-5)). In Proposition 3.5, we use the reduction mechanism of [Borel 2007; Mœglin and Waldspurger 1995] to obtain absolute convergence of the cuspidal Eisenstein series $E_f(s, g)$ from that of Borel Eisenstein series established in [Carbone et al. 2020].

Denote by ϖ_P the fundamental weight associated to α_P , and by ρ_M the Weyl vector of M . Let

$$W^\theta = \{w \in W : w^{-1}\alpha_i > 0, i \in \theta\}.$$

Then P is said to satisfy *property RD* if there exists a constant $D > 0$, such that for every nontrivial element $w \in W^\theta$ we have

$$\langle D\varpi_P + \rho_M, \alpha^\vee \rangle \leq 0$$

for any positive root α such that $w^{-1}\alpha < 0$. Property RD states that the coefficient of the simple root α_P grows faster than the coefficients of the simple roots in the subset θ . This property allows us to make use of the rapid decay of cusp forms on parabolic subgroups.

Now we state our main theorem.

Theorem 1.1. *Let f be an unramified cusp form on $(L \cap \Gamma) \backslash L$. If the maximal parabolic subgroup P satisfies property RD, then for any compact subset \mathfrak{S} of $A_{\mathcal{C}}$, there exists a measure zero subset S_0 of $(\Gamma \cap U) \backslash U\mathfrak{S}$ such that $E_f(s, g)$ is an entire function of $s \in \mathbb{C}$ for $g \in (\Gamma \cap U) \backslash U\mathfrak{S} - S_0K$.*

Here a measure zero set appears because absolute convergence is only established almost everywhere for the Eisenstein series in general. In the setting of everywhere convergence (as in [Carbone et al. 2020]), the measure zero set is not needed. See Theorem 4.8 for more precise statements.

The main idea of the proof is to exploit rapid decay of a cusp form on the maximal parabolic subgroup, guaranteed by property RD. This property may be considered as extraction of the essential properties of maximal parabolic subgroups used in the earlier results on affine Kac–Moody groups [Garland et al. 2017] and on rank 2 hyperbolic groups [Carbone et al. 2017].

In Section 5 we show that a large class of Kac–Moody groups have parabolic subgroups satisfying property RD, including the Kac–Moody group G associated with the Feingold–Frenkel rank 3 hyperbolic Kac–Moody algebra [Feingold and Frenkel 1983]. We note that the corresponding Kac–Moody group did not satisfy the conditions in [Carbone et al. 2020] that allowed the authors to prove convergence of Eisenstein series.

It would be an interesting question for future investigation to characterize the full class of Kac–Moody groups that admit parabolic subgroups satisfying property RD.

2. Kac–Moody groups and Borel Eisenstein series

In this section we recall the construction of Kac–Moody groups and then state the convergence results for Borel Eisenstein series proven in [Carbone et al. 2020] and used in this work. We try to keep the exposition self-contained as much as possible, and refer other details to [Carbone et al. 2020].

2A. Kac–Moody groups. Let $I = \{1, 2, \dots, r\}$, $A = (a_{ij})_{i,j \in I}$ be an $r \times r$ symmetrizable generalized Cartan matrix, and $(\mathfrak{h}_{\mathbb{C}}, \Delta, \Delta^{\vee})$ be a realization of A , where $\Delta = \{\alpha_1, \dots, \alpha_r\} \subset \mathfrak{h}_{\mathbb{C}}^*$ and $\Delta^{\vee} = \{\alpha_1^{\vee}, \dots, \alpha_r^{\vee}\} \subset \mathfrak{h}_{\mathbb{C}}$ are the set of simple roots and set of simple coroots, respectively.

Following [Carbone et al. 2020], we assume that A is nonsingular, so that $\mathfrak{h}_{\mathbb{C}}$ and $\mathfrak{h}_{\mathbb{C}}^*$ are spanned by the simple roots α_i and simple coroots α_i^{\vee} , respectively. Recall that $\langle \alpha_j, \alpha_i^{\vee} \rangle = a_{ij}$ for $i, j \in I$, where $\langle \cdot, \cdot \rangle$ is the natural pairing between $\mathfrak{h}_{\mathbb{C}}^*$ and $\mathfrak{h}_{\mathbb{C}}$. Denote the fundamental weights by $\varpi_i \in \mathfrak{h}_{\mathbb{C}}^*$, $i \in I$, which form the basis of $\mathfrak{h}_{\mathbb{C}}^*$ dual to the α_i^{\vee} .

Let $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g}_{\mathbb{C}}(A)$ be the Kac–Moody algebra associated to $(\mathfrak{h}, \Delta, \Delta^{\vee})$, which we assume to be infinite-dimensional throughout this paper. We denote by Φ the set of roots of $\mathfrak{g}_{\mathbb{C}}$ and have $\Phi = \Phi_+ \sqcup \Phi_-$, where Φ_+ (resp. Φ_-) is the set of positive (resp. negative) roots corresponding to the choice of Δ . Let $w_i := w_{\alpha_i}$ denote the simple Weyl reflection associated to the simple root α_i . The w_i for $i \in I$ generate the Weyl group W of $\mathfrak{g}_{\mathbb{C}}$.

Let \mathfrak{g} be the real Lie subalgebra generated by the Chevalley generators e_i and f_i for $i \in I$, so that $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$. As is standard (see [Carbone and Garland 2003, §4]), we can define the \mathbb{Z} -form $\mathcal{U}_{\mathbb{Z}}$ of the universal enveloping algebra $\mathcal{U}_{\mathbb{C}}$ of \mathfrak{g} using the Chevalley generators.

We now review the representation theoretic Kac–Moody groups G_F associated to \mathfrak{g} and a field F of characteristic zero, following [Carbone and Garland 2003]. Let (π, V) denote the unique irreducible highest weight module for \mathfrak{g} corresponding to a choice of some regular dominant integral weight, and fix

a nonzero highest weight vector $v \in V$. Then we have a lattice

$$V_{\mathbb{Z}} = \mathcal{U}_{\mathbb{Z}} \cdot v,$$

and we put $V_F = F \otimes_{\mathbb{Z}} V_{\mathbb{Z}}$. Since V is integrable, it makes sense to define for $s, t \in F$ and $i \in I$ the operators

$$u_{\alpha_i}(s) = \exp(\pi(se_i)) \quad \text{and} \quad u_{-\alpha_i}(t) = \exp(\pi(tf_i)),$$

which act locally finitely on V_F and thus define elements of $\text{Aut}(V_F)$.

For $t \in F^{\times}$ and $i \in I$ we set

$$w_{\alpha_i}(t) = u_{\alpha_i}(t)u_{-\alpha_i}(-t^{-1})u_{\alpha_i}(t)$$

and define

$$h_{\alpha_i}(t) = w_{\alpha_i}(t)w_{\alpha_i}(1)^{-1}.$$

Each simple root α_j defines a character on $\{h_{\alpha_i}(t) : t \in F^{\times}\}$ by

$$h_{\alpha_i}(t)^{\alpha_j} = t^{(\alpha_j, \alpha_i^{\vee})}. \quad (2-1)$$

The subgroup $\langle w_{\alpha_i}(1) : i \in I \rangle$ of $\text{Aut}(V_F)$ contains a full set of Weyl group representatives. For a real root α , choose once and for all an expression $\alpha = w_{\beta_1} \cdots w_{\beta_\ell} \alpha_i$ for some $i \in I$ and $\beta_1, \dots, \beta_\ell \in \Delta$, and define the corresponding one-parameter subgroup

$$u_{\alpha}(s) = wu_{\alpha_i}(s)w^{-1} \in \text{Aut}(V_F) \quad (s \in F),$$

where $w = w_{\beta_1}(1) \cdots w_{\beta_\ell}(1)$.

Define

$$G_F^0 = \langle u_{\alpha_i}(s), u_{-\alpha_i}(t) : s, t \in F, i \in I \rangle \subset \text{Aut}(V_F).$$

Following [Carbone and Garland 2003, §5], we choose a coherently ordered basis Ψ of $V_{\mathbb{Z}}$, and denote by B_F^0 the subgroup of G_F^0 consisting of elements which act upper-triangularly with respect to Ψ . The basis Ψ induces certain topologies on B_F^0 and G_F^0 , and we let B_F and G_F denote their completions respectively. When the field F is unspecified, B should be interpreted as $B_{\mathbb{R}}$.

The following subgroups of G_F are of particular importance to us:

- $A_F = \langle h_{\alpha_i}(s) : s \in F^{\times}, i \in I \rangle$.
- $U_F \subset B_F$, which is defined exactly as B_F , but with the additional stipulation that elements act unipotently upper-triangularly with respect to Ψ . It contains all subgroups parametrized by the $u_{\alpha}(\cdot)$, where $\alpha \in \Phi_+$ is a real root. Then $B_F = U_F A_F = A_F U_F$.

When no subscript is given, A and U should be interpreted as $A_{\mathbb{R}}$ and $U_{\mathbb{R}}$. We also have the following subgroups specific to the situation $F = \mathbb{R}$:

- K is the subgroup of $G_{\mathbb{R}}$ generated by all $\exp(t(e_i - f_i))$, $t \in \mathbb{R}$ and $i \in I$ ([Kac and Peterson 1985]).
- $A^+ = \langle h_{\alpha_i}(s) : s \in \mathbb{R}_{>0}, i \in I \rangle$. In fact, $(\mathbb{R}_{>0})^r$ can be identified with A^+ via the isomorphism $(x_1, \dots, x_r) \mapsto h_{\alpha_1}(x_1) \cdots h_{\alpha_r}(x_r)$, under which A^+ has the Haar measure da corresponding to $\prod_{i=1}^r \frac{dx_i}{x_i}$.

Theorem 2.1 [De Medts et al. 2009]. *We have the Iwasawa decomposition*

$$G_{\mathbb{R}} = UA^+K \tag{2-2}$$

with uniqueness of expression.

Remark 2.2. In [De Medts et al. 2009], twin BN-pairs are used to prove the Iwasawa decomposition for the group G^0 . The Iwasawa decomposition for $G_{\mathbb{R}}$ follows, since

$$G_{\mathbb{R}} = \bigcup_{w \in W} BwB = \bigcup_{w \in W} BwU_w = BG^0 = BB^0K = BK = UA^+K,$$

where $U_w := U \cap w^{-1}B_-^0w \subset G^0$ with the opposite Borel denoted by B_-^0 .

Let $u(g)$, $a(g)$, and $k(g)$ denote the projections from $G_{\mathbb{R}}$ onto each of the respective factors in (2-2). We define the discrete group $\Gamma := G_{\mathbb{R}} \cap \text{Aut}(V_{\mathbb{Z}}) = \{\gamma \in G_{\mathbb{R}} : \gamma \cdot V_{\mathbb{Z}} = V_{\mathbb{Z}}\}$. As in [Garland 2004], it can be shown that $(\Gamma \cap U) \backslash U$ is the projective limit of a projective family of finite-dimensional compact nil-manifolds and thus admits a projective limit measure du which is a right U -invariant probability measure.

2B. Borel Eisenstein series. Using the identification of A^+ with $(\mathbb{R}_{>0})^r$, each element of $\mathfrak{h}_{\mathbb{C}}^*$ gives rise to a quasicharacter of A^+ . Let $\lambda \in \mathfrak{h}_{\mathbb{C}}^*$ and let $\rho \in \mathfrak{h}_{\mathbb{C}}^*$ be the Weyl vector, which is characterized by $\langle \rho, \alpha_i^\vee \rangle = 1$, $i \in I$. We set

$$\begin{aligned} \Phi_\lambda : G_{\mathbb{R}} &\rightarrow \mathbb{C}^\times, \\ g &\mapsto a(g)^{\lambda+\rho}, \end{aligned} \tag{2-3}$$

which is well-defined by the uniqueness of the Iwasawa decomposition.

Let B and Γ be as defined in Section 2A. Define the Borel Eisenstein series on $G_{\mathbb{R}}$ to be the infinite formal sum

$$E_\lambda(g) := \sum_{\gamma \in (\Gamma \cap B) \backslash \Gamma} \Phi_\lambda(\gamma g).$$

We define for all $g \in G_{\mathbb{R}}$ the so-called “upper triangular” constant term¹

$$E_\lambda^\sharp(g) := \int_{(\Gamma \cap U) \backslash U} E_\lambda(ug) du,$$

which is left U -invariant and right K -invariant. It is immediate from the definition that $E_\lambda^\sharp(g)$ is determined by the A^+ -component $a(g)$ of g in the Iwasawa decomposition.

We now state the convergence results for the constant term $E_\lambda^\sharp(g)$ as well as the Eisenstein series $E_\lambda(g)$ itself obtained in [Carbone et al. 2020]. We will use these results later in this paper.

Let $\mathcal{C} \subset \mathfrak{h}$ be the open fundamental chamber

$$\mathcal{C} = \{x \in \mathfrak{h} : \langle \alpha_i, x \rangle > 0, i \in I\}.$$

¹This is to be distinguished from the proposal in [Braverman and Kazhdan 2013] to consider constant terms in uncompleted, or so-called “lower triangular”, parabolics.

Let \mathfrak{C} denote the interior of the Tits cone $\bigcup_{w \in W} w\bar{\mathcal{C}}$ corresponding to \mathcal{C} [Kac 1990, §3.12]. Using the exponential map $\exp : \mathfrak{h} \rightarrow A^+$, set $A_{\mathcal{C}} = \exp \mathcal{C}$ and $A_{\mathfrak{C}} = \exp \mathfrak{C}$. Let

$$\mathcal{C}^* = \{\lambda \in \mathfrak{h}^* : \langle \lambda, \alpha_i^\vee \rangle > 0, i \in I\}. \tag{2-4}$$

Based on a crucial lemma due to Looijenga [1980] together with some elementary estimate on the Riemann ζ -function, the following theorem is proven in [Carbone et al. 2020].

Theorem 2.3. *If $\lambda \in \mathfrak{h}_{\mathbb{C}}^*$ satisfies $\operatorname{Re}(\lambda - \rho) \in \mathcal{C}^*$, then $E_{\lambda}^{\sharp}(g)$ converges absolutely for $g \in UA_{\mathfrak{C}}K$, and in fact uniformly for $a(g)$ lying in any fixed compact subset of $A_{\mathfrak{C}}$.*

Consequently, a simple application of Tonelli’s theorem as in [Garland 2004, §9] yields the following.

Corollary 2.4 [Carbone et al. 2020]. *For $\lambda \in \mathfrak{h}_{\mathbb{C}}^*$ with $\operatorname{Re}(\lambda - \rho) \in \mathcal{C}^*$ and any compact subset \mathfrak{S} of $A_{\mathfrak{C}}$, there exists a measure zero subset S_0 of $(\Gamma \cap U) \backslash U\mathfrak{S}$ such that the series $E_{\lambda}(g)$ converges absolutely for $g \in (\Gamma \cap U) \backslash U\mathfrak{S}K - S_0K$.*

Moreover, the everywhere convergence of $E_{\lambda}(g)$ over $\Gamma UA_{\mathfrak{C}}K$ was established under the following combinatorial property, by exploring the idea that the Borel Eisenstein series can be nearly bounded by its constant term. We set

$$\Phi_w := \Phi_+ \cap w^{-1}\Phi_-, \quad w \in W. \tag{2-5}$$

The following property was introduced in [Carbone et al. 2020] and will be used in Theorem 4.8.

Property 2.5. Every nontrivial $w \in W$ can be written as $w = vw_{\beta}$, where β is a positive simple root, $\ell(v) < \ell(w)$, and $\alpha - \beta$ is never a real root for any $\alpha \in \Phi_v$.

In [Carbone et al. 2020] it is shown that Property 2.5 holds when the Cartan matrix $A = (a_{ij})$ is symmetric with $|a_{ij}| \geq 2$ for all i, j ; for rank 2 hyperbolic groups, the condition $|a_{ij}| \geq 2$ is sufficient and the matrix A does not need to be symmetric. The main result of that paper is the following.

Theorem 2.6 [Carbone et al. 2020]. *Assume that $\lambda \in \mathfrak{h}_{\mathbb{C}}^*$ satisfies $\operatorname{Re}(\lambda - \rho) \in \mathcal{C}^*$, and that Property 2.5 holds. Then the Kac–Moody Eisenstein series $E_{\lambda}(g)$ converges absolutely for $g \in \Gamma UA_{\mathfrak{C}}K$.*

3. Cuspidal Kac–Moody Eisenstein series

3A. Definition of Eisenstein series. For any subset $\theta \subset I$, denote by W_{θ} the subgroup of W generated by the reflections w_i for $i \in \theta$. Assume that we choose θ such that W_{θ} is a finite group. Define the parabolic subgroup of $G_{\mathbb{R}}$ associated to θ by

$$P_{\theta} := BW_{\theta}B,$$

where each $w \in W_{\theta}$ is identified with one of its representatives in $G_{\mathbb{R}}$ as in Section 2A. For the rest of this section, we will fix θ and write $P = P_{\theta}$.

Let N be the pro-unipotent radical of P , and M be the subgroup of P generated by A and $u_{\pm\alpha_i}(t)$, $i \in \theta$, $t \in \mathbb{R}$. Then M is a finite-dimensional real reductive group. By [Kumar 2002, Lemma 6.1.13], we have the Levi decomposition $P = M \ltimes N$.

We shall introduce certain subgroups and explain some structures of M . Let L be the subgroup of M generated by $u_{\pm\alpha_i}(t)$, $i \in \theta$, $t \in \mathbb{R}$. Then L is a finite-dimensional semisimple group. Let $A_1 := A \cap L$ be the subgroup of A generated by $h_{\alpha_i}(t)$, $t \in \mathbb{R}^\times$, $i \in \theta$, which is a maximal split torus of L . Set

$$A_1^+ = \langle h_{\alpha_i}(t) : t \in \mathbb{R}_{>0}, i \in \theta \rangle \cong A_1 / (A_1 \cap K).$$

We also introduce

$$H := \{a \in A : a^{\alpha_i} = \pm 1, i \in \theta\},$$

and let H^+ be the identity component of H , so that $H^+ \cong H / (H \cap K)$.

Lemma 3.1. *We have $A \cong A_1^+ \times H$ and $A^+ \cong A_1^+ \times H^+$.*

Proof. The second isomorphism follows from taking the identity component of the first. Let us prove the first one. By the assumption on θ , the natural pairing between $\bigoplus_{i \in \theta} \mathbb{Z}\alpha_i$ and $\bigoplus_{i \in \theta} \mathbb{Z}\alpha_i^\vee$ is nondegenerate. From this it follows easily that $A_1^+ \cap H = \{1\}$. For the same reason, for any $a \in A$, there exists $a_1 \in A_1^+$ such that $a_1^{\alpha_i} = |a^{\alpha_i}|$ for all $i \in \theta$. Then clearly $aa_1^{-1} \in H$, which proves that $A = A_1^+H$ and therefore $A \cong A_1^+ \times H$. \square

Remark 3.2. By Lemma 3.1 it is clear that $M = LH$, and we can say a bit more about this decomposition. Note that H^+ is the identity component of the center of M . By [Borel 2007, §4.3.1], we have

$$M \cong {}^\circ M \times H^+,$$

where

$${}^\circ M := \bigcap_{\chi \in X(M)} \ker|\chi|.$$

Here $X(M)$ denotes the lattice of algebraic characters of M defined over \mathbb{R} . Since L is semisimple, we have $L \subset {}^\circ M$, hence $L \cap H \subset {}^\circ M \cap H$, which is a finite group contained in K .

It will be convenient to rewrite $a(g)$ as

$$\text{Iw}_{A^+} : G_{\mathbb{R}} \rightarrow A^+, \quad g \mapsto a(g)$$

from the Iwasawa decomposition $G_{\mathbb{R}} = UA^+K$ in (2-2). The parabolic subgroup P gives rise to the decomposition $G_{\mathbb{R}} = NMK$, which unlike the Iwasawa decomposition (2-2) is not unique since $M \cap K$ is nontrivial. However in Lemma 3.3 we will show that the projection

$$\text{Iw}_M : G_{\mathbb{R}} \rightarrow M / (M \cap K)$$

is well-defined. Although $M = LH$ is in general not a direct product (see Remark 3.2), we will also deduce from the fact $L \cap H \subset K$ that the maps

$$\text{Iw}_L : G_{\mathbb{R}} \rightarrow L / (L \cap K) \quad \text{and} \quad \text{Iw}_{H^+} : G_{\mathbb{R}} \rightarrow H^+,$$

which are induced by the projections onto the factors L and H of M , respectively, are well-defined.

Similarly, using the Iwasawa decomposition for L , we may define the map

$$\text{Iw}_{A_1^+} : L/(L \cap K) \rightarrow A_1^+.$$

Lemma 3.3. *The Iwasawa maps Iw_M , Iw_L and Iw_{H^+} are well-defined.*

Proof. Well-definedness of Iw_M : Assume that $nmk = m'$, where $n \in N$, $m, m' \in M$ and $k \in K$. We need to show that $m' \in mK$, or that $n = 1$. Apply the Iwasawa decomposition of M to write $m = u_M a_M k_M$ and $m' = u'_M a'_M k'_M$, where $u_M, u'_M \in U_M := U \cap M$, $a_M, a'_M \in A^+$ and $k_M, k'_M \in K \cap M$. By the uniqueness of the Iwasawa decomposition $G = UA^+K$, we have in particular that

$$nu_M = u'_M,$$

which forces $n = 1$ and $u_M = u'_M$ because $U = U_M \rtimes N$. Hence $m' = mk$, which implies that Iw_M is well-defined.

Well-definedness of Iw_L and Iw_{H^+} : In Remark 3.2 we have shown that $M = LH$ and $L \cap H \subset K$. Since we already proved that Iw_M is well-defined, we now only need to consider

$$lh = l'h' \in M,$$

where $l, l' \in L$ and $h, h' \in H$. Then $l^{-1}l' = h(h')^{-1} = (h')^{-1}h \in L \cap H \subset K$; hence $l' \in lK$, $h \in h'K$, which prove that Iw_L and Iw_{H^+} are both well-defined. \square

We have the following decomposition of the Iwasawa map Iw_{A^+} .

Lemma 3.4. *We have*

$$\text{Iw}_{A^+} = (\text{Iw}_{A_1^+} \circ \text{Iw}_L) \times \text{Iw}_{H^+} : G_{\mathbb{R}} \rightarrow A^+ \cong A_1^+ \times H^+. \quad (3-1)$$

Proof. We can verify (3-1) by tracking the definitions. Assume that

$$g = nmk, \quad m = lh,$$

where each factor belongs to the subgroup denoted by the corresponding capital letter. Applying the Iwasawa decomposition of L , we can write

$$l = ua_1k_L,$$

where $u \in U_M$, $a_1 \in A_1^+$ and $k_L \in K \cap L$. We can also write $h = h^+k_H$, where $h^+ \in H^+$ and $k_H \in K \cap H$. Since H^+ commutes with L , we have

$$g = (nu)(a_1h^+)(k_Lk_Hk),$$

which is the Iwasawa decomposition according to $G_{\mathbb{R}} = UA^+K$. By the definitions of various Iwasawa maps, we have

$$\text{Iw}_{A^+} = a_1h^+ = \text{Iw}_{A_1^+}(lK_L) \cdot \text{Iw}_{H^+}(g) = \text{Iw}_{A_1^+}(\text{Iw}_L(g)) \cdot \text{Iw}_{H^+}(g),$$

which proves (3-1). \square

Consider $\lambda_P = \sum_{i \notin \theta} s_i \varpi_i \in \mathfrak{h}_{\mathbb{C}}^*$, $s_i \in \mathbb{C}$, where ϖ_i are the fundamental weights. As before, let $\rho \in \mathfrak{h}_{\mathbb{C}}^*$ be the Weyl vector, i.e., $\langle \rho, \alpha_i^\vee \rangle = 1$, $i \in I$. Likewise, let ρ_M be half the sum of the positive roots of M , and set $\rho_P := \rho - \rho_M$. Since λ_P is clearly trivial on $A_1 = A \cap L$, the decomposition (3-1) implies that

$$\mathrm{Iw}_{A^+}(\cdot)^{\lambda_P} = \mathrm{Iw}_{H^+}(\cdot)^{\lambda_P}. \quad (3-2)$$

Similarly, every root β of L is trivial on H^+ ; hence

$$\mathrm{Iw}_{A^+}(\cdot)^\beta = (\mathrm{Iw}_{A_1^+} \circ \mathrm{Iw}_L(\cdot))^\beta. \quad (3-3)$$

For any $\lambda_P = \sum_{i \notin \theta} s_i \varpi_i \in \mathfrak{h}_{\mathbb{C}}^*$, define the auxiliary Eisenstein series

$$E(\lambda_P, g) := \sum_{\gamma \in (\Gamma \cap P) \backslash \Gamma} \mathrm{Iw}_{H^+}(\gamma g)^{\lambda_P + \rho_P}, \quad g \in G_{\mathbb{R}}. \quad (3-4)$$

More generally, for a bounded function f defined on $M/M \cap K$, define the Eisenstein series $E_f(\lambda_P, g)$ by

$$E_f(\lambda_P, g) := \sum_{\gamma \in (\Gamma \cap P) \backslash \Gamma} \mathrm{Iw}_{H^+}(\gamma g)^{\lambda_P + \rho_P} f(\mathrm{Iw}_M(\gamma g)), \quad g \in G_{\mathbb{R}}. \quad (3-5)$$

Then (3-4) corresponds to the special case that $f \equiv 1$.

3B. Convergence by the reduction mechanism. The idea of a reduction mechanism is to bound general Eisenstein series by Borel ones. It was first suggested by Bernstein to Borel, and used by Garland [2011] for the affine case, inspired by some results in [Green et al. 2010]. Following this idea, we prove the absolute convergence of parabolic Eisenstein series $E_f(\lambda_P, g)$ in the Godement range. Similar proofs for the finite-dimensional case can be found in [Borel 2007, Proposition 12.6] and [Mœglin and Waldspurger 1995, Proposition II.1.5], which also use Borel Eisenstein series to bound parabolic ones, but the arguments are slightly different.

Proposition 3.5. *Let f be a bounded function on $M/M \cap K$. For $\lambda_P = \sum_{i \notin \theta} s_i \varpi_i \in \mathfrak{h}_{\mathbb{C}}^*$, with $\mathrm{Re}(s_i) > \langle \rho_P, \alpha_i^\vee \rangle$, $i \notin \theta$, and any compact subset \mathfrak{S} of $A_{\mathbb{C}}$, there exists a measure zero subset S_0 of $(\Gamma \cap U) \backslash U \mathfrak{S}$ such that the series $E_f(\lambda_P, g)$ converges absolutely for $g \in (\Gamma \cap U) \backslash U \mathfrak{S} K - S_0 K$. Moreover, S_0 can be chosen to work uniformly for all s_i , $i \notin \theta$, in right half-spaces of the form $\mathrm{Re}(s_i) > \sigma_i$, where $\sigma_i > \langle \rho_P, \alpha_i^\vee \rangle$ are fixed.*

Proof. Since the function f is assumed to be bounded, it suffices to prove the almost everywhere absolute convergence of the auxiliary Eisenstein series $E(\lambda_P, g)$, and we may assume that $s_i > \langle \rho_P, \alpha_i^\vee \rangle$ are real numbers for $i \notin \theta$. Let $\varepsilon > 0$ be sufficiently small such that

$$s_i > (1 + \varepsilon) \langle \rho_P, \alpha_i^\vee \rangle$$

for $i \notin \theta$. We put

$$\lambda = (1 + \varepsilon) \rho_M + \lambda_P.$$

Then $\lambda - \rho \in \mathcal{C}^*$, where \mathcal{C}^* is defined in (2-4). Indeed, $\langle \lambda, \alpha_i^\vee \rangle = 1 + \varepsilon$ for $i \in \theta$, and

$$\langle \lambda, \alpha_i^\vee \rangle = (1 + \varepsilon)\langle \rho_M, \alpha_i^\vee \rangle + s_i > (1 + \varepsilon)\langle \rho_M + \rho_P, \alpha_i^\vee \rangle = 1 + \varepsilon$$

for $i \notin \theta$.

Consider the Borel Eisenstein series

$$E_\lambda(g) = \sum_{\gamma \in (\Gamma \cap B) \backslash \Gamma} \Phi_\lambda(\gamma g) = \sum_{\gamma \in (\Gamma \cap P) \backslash \Gamma} \sum_{\delta \in (\Gamma \cap B) \backslash \Gamma \cap P} \Phi_\lambda(\delta \gamma g).$$

By the Iwasawa decomposition $G_{\mathbb{R}} = NMK$ and Lemma 3.6, the last inner sum is equal to

$$\sum_{\delta \in (\Gamma_M \cap B_M) \backslash \Gamma_M} \Phi_\lambda(\delta \cdot \text{Iw}_M(\gamma g)),$$

where $\Gamma_M := \Gamma \cap M$ and $B_M := B \cap M$. By our choice of λ , this can be further manipulated to give

$$\text{Iw}_{H^+}(\gamma g)^{\lambda_P + \rho_P} \sum_{\delta \in (\Gamma_M \cap B_M) \backslash \Gamma_M} \Phi_\lambda(\delta \cdot \text{Iw}_L(\gamma g)) = \text{Iw}_{H^+}(\gamma g)^{\lambda_P + \rho_P} \sum_{\delta \in (\Gamma_M \cap B_M) \backslash \Gamma_M} \text{Iw}_{A_1^+}(\delta \cdot \text{Iw}_L(\gamma g))^{\lambda + \rho}.$$

Since λ and ρ restrict to $(1 + \varepsilon)\rho_M$ and ρ_M on A_1^+ respectively, the last sum is a Borel Eisenstein series on M with spectral parameter $(1 + \varepsilon)\rho_M$, which is in the Godement range. Since the spectral parameter is real, the series is a sum of positive terms, and hence is never zero. Moreover, it has a positive lower bound. This fact should be well known, but it does not seem to be widely available in the literature. We supply a proof below for completeness.

Recall that a Siegel domain for M is of the form

$$\mathfrak{S}_{t,\omega} := \omega A_t K_M,$$

where ω is a compact subset of B_M , $K_M = K \cap M$ and

$$A_t := \{a \in A^+ : a^{\alpha_i} > t, i \in \theta\}$$

for $t > 0$. By classical reduction theory (see, e.g., [Mœglin and Waldspurger 1995, I.2.1]), we have $M = \Gamma_M \cdot \mathfrak{S}_{t,\omega}$ for ω sufficiently large and t sufficiently small. Write $m \in M$ accordingly as

$$m = \gamma x a k,$$

where $\gamma \in \Gamma_M$, $x \in \omega$, $a \in A_t$ and $k \in K_M$. The Borel Eisenstein series $E_{(1+\varepsilon)\rho_M}(\cdot)$ on M takes the value

$$\begin{aligned} E_{(1+\varepsilon)\rho_M}(m) &= E_{(1+\varepsilon)\rho_M}(xa) \geq \Phi_{(1+\varepsilon)\rho_M}(xa) = a^{(2+\varepsilon)\rho_M} \Phi_{(1+\varepsilon)\rho_M}(x) \\ &\geq \kappa := t^{(2+\varepsilon)C_M} \cdot h_\omega > 0, \end{aligned}$$

where C_M is the sum of the coefficients of α_i in ρ_M and h_ω is the minimum value of the positive continuous function $\Phi_{(1+\varepsilon)\rho_M}(\cdot)$ on the compact set ω .

With this lower bound established, it follows that

$$E(\lambda_P, g) = \sum_{\gamma \in (\Gamma \cap P) \backslash \Gamma} \text{Iw}_{H^+}(\gamma g)^{\lambda_P + \rho_P} \leq \kappa^{-1} E_\lambda(g).$$

The proposition follows from this comparison and Corollary 2.4. \square

Lemma 3.6. *If $\gamma \in \Gamma \cap P$ decomposes as $\gamma = nm$, where $n \in N$, $m \in M$, then $n, m \in \Gamma$.*

Proof. Recall from [Carbone and Garland 2003, §5] that a coherently ordered basis Ψ of $V_{\mathbb{Z}}$ consists of weight vectors. If $m \notin \Gamma$, then there exists $v_0 \in \Psi$ such that $m \cdot v_0 \notin V_{\mathbb{Z}}$. Assume that v_0 is of weight μ . Then there exists a finite subset $\Psi_0 \subset \Psi$ such that each $v \in \Psi_0$ is of weight $\mu + \alpha$ for some α in the root lattice of M , and that $m \cdot v_0$ is of the form

$$m \cdot v_0 = \sum_{v \in \Psi_0} c_v v \quad \text{with } c_v \notin \mathbb{Z} \text{ for some } v \in \Psi_0.$$

It is clear that $nm \cdot v_0$ is of the form

$$nm \cdot v_0 = m \cdot v_0 + \sum_{v \in \Psi \setminus \Psi_0} c_v v \notin V_{\mathbb{Z}},$$

which contradicts that $\gamma = nm \in \Gamma$. \square

4. Holomorphy of cuspidal Eisenstein series

In this section, we prove our main result. We keep the assumption that \mathfrak{g} is infinite-dimensional with nonsingular Cartan matrix A . Let $P = P_\theta$ be a maximal parabolic subgroup of $G_{\mathbb{R}}$ with fixed Levi decomposition and associated finite-dimensional split reductive Levi subgroup M as in Section 3. We set $\{i_P\} := I \setminus \theta$ and recall that $\alpha_P = \alpha_{i_P}$, and ϖ_P is the fundamental weight for α_P . Let W^θ be the set of minimal length representatives of $W_\theta \backslash W$, i.e.,

$$W^\theta := \{w \in W : w^{-1}\alpha_i > 0, i \in \theta\}.$$

Definition 4.1. A maximal parabolic subgroup $P = P_\theta$ whose Levi subgroup M is a finite-dimensional split reductive group, is said to satisfy *property RD* if there exists a constant $D > 0$, such that for every nontrivial element $w \in W^\theta$ we have $\langle D\varpi_P + \rho_M, \alpha^\vee \rangle \leq 0$ for any $\alpha \in \Phi_{w^{-1}}$.

Remark 4.2. Note that $\langle \varpi_P, \alpha^\vee \rangle > 0$ for any $\alpha \in \Phi_{w^{-1}}$ with $w \in W^\theta$. Hence if there exists D that satisfies the condition in Definition 4.1, then any positive constant smaller than D will also satisfy the condition. Actually, one can see that P_θ has property RD if and only if for every nontrivial element $w \in W^\theta$ we have $\langle \rho_M, \alpha^\vee \rangle < 0$ for any $\alpha \in \Phi_{w^{-1}}$.

The fact $L \cap H \subset K$ allows us to view a function f on $L/L \cap K$ as defined on $M/M \cap K$. In particular, an unramified cusp form f on L can be regarded as a bounded function on $M/M \cap K$, hence one may define the Eisenstein series $E_f(\lambda_P, g)$ as in Section 3. In the sequel we shall adopt this interpretation without further remark. Moreover, since $\lambda_P = s\varpi_P$ for some $s \in \mathbb{C}$, we write $E_f(s, g) = E_f(\lambda_P, g)$ in the rest of the paper.

We state the main theorem again.

Theorem 4.3 (Theorem 1.1). *Let f be an unramified cusp form on $L \cap \Gamma \backslash L$. If the maximal parabolic subgroup P satisfies property RD, then for any compact subset \mathfrak{S} of $A_{\mathbb{C}}$, there exists a measure zero subset S_0 of $(\Gamma \cap U) \backslash U \mathfrak{S}$ such that $E_f(s, g)$ is an entire function of $s \in \mathbb{C}$ for $g \in (\Gamma \cap U) \backslash U \mathfrak{S} K - S_0 K$.*

The rest of this section is devoted to the proof of Theorem 4.3. We shall follow the strategy of [Garland et al. 2017] and [Carbone et al. 2017] with property RD at hand. We prove an inequality in Lemma 4.7 which is crucial in our proof of the main theorem. This inequality enables us to deduce the entirety of Eisenstein series from the rapid decay of cusp forms. Along the way, we note some interesting properties of the infinite-dimensional Kac–Moody algebras that are considered here.

The following lemma is essentially the same as [Garland et al. 2017, Lemma 3.2]. Write $x^y = yxy^{-1}$ for $x, y \in G_{\mathbb{R}}$, and set $U_w = \prod_{\alpha \in \Phi_w} U_{\alpha}$ for $w \in W$ where U_{α} is the one-parameter subgroup corresponding to α .

Lemma 4.4. *Let $w \in W$. If $\gamma \in \Gamma \cap BwB$, then*

$$Iw_{A^+}(\gamma g) = Iw_{A^+}(g)^w \cdot Iw_{A^+}(wu_w)$$

for some $u_w \in U_w$ depending on γ and g .

We also recall that one has, for $u_w \in U_w$,

$$\log(Iw_{A^+}(wu_w)) = \sum_{\alpha \in \Phi_{w^{-1}}} c_{\alpha} \alpha^{\vee} \quad \text{with } c_{\alpha} \leq 0 \tag{4-1}$$

by [Garland 2007b, Lemma 6.1] or [Garland et al. 2017, (3.25)].

To utilize property RD, we need the property of the coefficient of α_P in the fundamental weight ϖ_P , given by the following lemma.

Lemma 4.5. *The coefficient n_{i_P} of α_P in the expansion $\varpi_P = \sum_{i \in I} n_i \alpha_i$ is negative.*

Proof. By relabeling the simple roots, we may assume that $\alpha_P = \alpha_r$ and the upper-left $(r-1) \times (r-1)$ corner of A is the Cartan matrix of the root system Φ_M of M . Let $D = \text{diag}(d_1, \dots, d_r)$ be a symmetrizer of A and write

$$DA = \begin{pmatrix} U & \beta \\ \beta^T & c \end{pmatrix}, \quad (DA)^{-1} = \begin{pmatrix} V & \gamma \\ \gamma^T & d \end{pmatrix},$$

where U, V are $(r-1) \times (r-1)$ matrices and c, d are scalars. Then $U = U^T$ is positive definite because Φ_M is of finite type. If $\gamma^T = (\gamma_1, \dots, \gamma_{r-1})$, then

$$\varpi_r = \sum_{i=1}^{r-1} d_r \gamma_i \alpha_i + d_r d \alpha_r.$$

Hence we only need to show that $d < 0$.

The matrix DA has the same signature as

$$\begin{pmatrix} U & 0 \\ 0 & c - \beta^T U^{-1} \beta \end{pmatrix}.$$

Since we assume that \mathfrak{g} is of infinite type and A is nonsingular, we must have $c - \beta^T U^{-1} \beta < 0$. We have the relations

$$U\gamma + \beta d = 0, \quad \beta^T \gamma + cd = 1.$$

It follows that

$$\beta^T \gamma + \beta^T U^{-1} \beta d = \beta^T U^{-1} (U\gamma + \beta d) = 0,$$

which implies that $(c - \beta^T U^{-1} \beta)d = 1$. Hence $d < 0$ as desired. \square

As a corollary, we obtain the following result.

Lemma 4.6. *For $D > 0$ sufficiently small, $w^{-1}(D\varpi_P + \rho_M)$ is a positive linear combination of simple roots for any nontrivial $w \in W^\theta$.*

Proof. Recall that ρ_M is the half sum of the positive roots of M . Write $\varpi_P = \sum_{i \in I} n_i \alpha_i$ as in Lemma 4.5. Then we have

$$D\varpi_P + \rho_M = \sum_{i \in \theta} m_i \alpha_i + m_{i_P} \alpha_P,$$

where

$$\sum_{i \in \theta} m_i \alpha_i = \rho_M + D \sum_{i \in \theta} n_i \alpha_i,$$

and $m_{i_P} = Dn_{i_P} < 0$ by Lemma 4.5. Clearly for $D > 0$ sufficiently small we have $m_i > 0$ for $i \in \theta$. Consider a nontrivial element $w \in W^\theta$. Since $w^{-1} \alpha_i > 0$ for $i \in \theta$, we must have $w^{-1} \alpha_P < 0$. The lemma follows immediately. \square

With the above results, we can now state and prove the following inequality.

Lemma 4.7. *If P satisfies property RD, then for $D > 0$ sufficiently small,*

$$\mathrm{Iw}_{A^+}(\gamma g)^{-\rho_M} \leq \mathrm{Iw}_{A^+}(\gamma g)^{D\varpi_P}$$

for any $g \in UA_C K$, $w \in W^\theta$ and $\gamma \in \Gamma \cap BwB$.

Proof. From Definition 4.1, there exists $D > 0$ such that

$$\langle D\varpi_P + \rho_M, \alpha^\vee \rangle \leq 0 \quad \text{for } \alpha \in \Phi_{w^{-1}}. \quad (4-2)$$

By Remark 4.2, we may shrink the constant D , if necessary, and assume that the property in Lemma 4.6 holds with this constant D . By Lemma 4.4 and (4-1), in order to prove Lemma 4.7, it suffices to establish its logarithmic form

$$\langle w^{-1}(D\varpi_P + \rho_M), \log(\mathrm{Iw}_{A^+}(g)) \rangle + \sum_{\alpha \in \Phi_{w^{-1}}} c_\alpha \langle D\varpi_P + \rho_M, \alpha^\vee \rangle \geq 0, \quad (4-3)$$

with $c_\alpha \leq 0$, $\alpha \in \Phi_{w^{-1}}$. The first term is positive by Lemma 4.6 and the assumption that $\log(\mathrm{Iw}_{A^+}(g)) \in \mathcal{C}$. The summation in (4-3) is nonnegative by property RD (4-2). Hence (4-3) holds. \square

Proof of Theorem 4.3. Fix any real number $s_0 > \langle \rho_P, \alpha_P^\vee \rangle$, and assume that $\text{Re}(s) < s_0$. By the rapid decay of cuspidal automorphic forms (e.g., [Miller and Schmid 2012]), for any $n > 0$ there exists a constant $C_1 > 0$ depending on n such that

$$|f(g)| \leq C_1 \text{Iw}_{A_1^+}(g)^{-n\rho_M}. \tag{4-4}$$

We take

$$n := \frac{s_0 - \text{Re}(s)}{D} > 0,$$

with D the constant in Lemma 4.7.

Recall that we have the Bruhat decomposition ([Carbone and Garland 2003])

$$G_{\mathbb{R}} = \bigcup_{w \in W^\theta} PwB.$$

We claim that for any $\gamma \in (\Gamma \cap P) \setminus (\Gamma \cap PwB)$ with $w \in W^\theta$, we can choose a representative of γ which lies in $\Gamma \cap BwB$. Recall the Levi decomposition $P = MN$. By the Iwasawa decomposition over \mathbb{Q} , we have a representative of γ of the form

$$\tilde{\gamma} = mnwb \in \Gamma \cap PwB,$$

where $m \in M_{\mathbb{Q}}$, $n \in N_{\mathbb{Q}}$ and $b \in B_{\mathbb{Q}}$. Since M is finite-dimensional, we have $M_{\mathbb{Q}} = \Gamma_M B_{M,\mathbb{Q}}$ (see [Godement 1964]), where $\Gamma_M := \Gamma \cap M$ and $B_{M,\mathbb{Q}} := B_{\mathbb{Q}} \cap M$. Write $m = \gamma_M b_M$ accordingly, where $\gamma_M \in \Gamma_M$ and $b_M \in B_{M,\mathbb{Q}}$. Then

$$\gamma_M^{-1} \tilde{\gamma} = b_M n w b$$

is a representative of γ , which clearly lies in $\Gamma \cap BwB$.

With the claim established, it follows from (3-2), (3-3), Lemma 4.7 and (4-4) that

$$\begin{aligned} |\text{Iw}_{H^+}(\gamma g)^{s\varpi_P} f(\text{Iw}_M(\gamma g))| &\leq C_1 \text{Iw}_{H^+}(\gamma g)^{(\text{Re } s)\varpi_P} \text{Iw}_{A_1^+} \circ \text{Iw}_L(\gamma g)^{-n\rho_M} \\ &\leq C_1 \text{Iw}_{H^+}(\gamma g)^{(\text{Re } s)\varpi_P} \text{Iw}_{H^+}(\gamma g)^{nD\varpi_P} \\ &= C_1 \text{Iw}_{H^+}(\gamma g)^{s_0\varpi_P}. \end{aligned}$$

Note that the constants C_1 and D are independent of w and that the estimation works for any representative $\gamma \in \Gamma \cap BwB$. Multiplying by $\text{Iw}_{H^+}(\gamma g)^{\rho_P}$ and taking the summation over γ , it follows that $E_f(s, g)$ is bounded by $C_1 E(s_0, g)$. Hence by Proposition 3.5 the series $E_f(s, g)$ is absolutely convergent on $U\mathfrak{S}K$ off S_0K , with S_0 a measure zero subset of $(\Gamma \cap U) \setminus U\mathfrak{S}$. Note that this subset S_0 depends on s_0 . To conclude, we may take a sequence of real numbers $s_0^{(i)} > \langle \rho_P, \alpha_P^\vee \rangle$ which goes to infinity, and take the countable union of the corresponding measure zero subsets $S_0^{(i)}$. \square

Assuming Property 2.5, we have everywhere convergence of cuspidal Eisenstein series over the full Tits cone by the reduction mechanism. Thus in this case Proposition 3.5 and Theorem 4.3 can be strengthened. We summarize this as the following result.

Theorem 4.8. *Assume Property 2.5, and let $P = MN$ be a maximal parabolic subgroup of $G_{\mathbb{R}}$ with finite-dimensional Levi subgroup M . Then the following hold.*

- (1) *For a bounded function f on $M \cap \Gamma \backslash M / M \cap K$, the Eisenstein series $E_f(s, g)$ converges absolutely for $g \in \Gamma U A_{\mathbb{C}} K$ and $\operatorname{Re}(s) > 1$.*
- (2) *If P satisfies property RD and f is an unramified cusp form on $L \cap \Gamma \backslash L$, then the Eisenstein series $E_f(s, g)$ is an entire function of $s \in \mathbb{C}$ for $g \in \Gamma U A_{\mathbb{C}} K$.*

5. Parabolic subgroups satisfying property RD

In this section we investigate a sufficient condition which implies that every maximal parabolic subgroup of $G_{\mathbb{R}}$ with finite-dimensional Levi subgroup satisfies property RD, and then discuss some examples, including the rank 3 Feingold–Frenkel hyperbolic algebra \mathcal{F} .

We keep the notation of the previous section. In particular, $P = P_{\theta}$ stands for a maximal parabolic subgroup of $G_{\mathbb{R}}$, whose Levi subgroup M is a finite-dimensional split reductive group.

Proposition 5.1. *For any $w \in W$ and $\alpha \in \Phi_{w^{-1}}$, suppose that we have $\langle \alpha_i, \alpha^{\vee} \rangle \leq 0$ for any α_i such that $w^{-1}\alpha_i > 0$. Then every maximal parabolic subgroup P with finite-dimensional Levi subgroup satisfies property RD.*

Proof. As in the proof of Lemma 4.6, for small $D > 0$ we have

$$D\varpi_P + \rho_M = \sum_{i \in \theta} m_i \alpha_i + m_{i_P} \alpha_P$$

where $m_i > 0$ for $i \in \theta$ and $m_{i_P} < 0$. Take $w \in W^{\theta}$, $w \neq \operatorname{id}$, and $\alpha \in \Phi_{w^{-1}}$. Then by assumption

$$\langle \alpha_i, \alpha^{\vee} \rangle \leq 0 \quad \text{for } i \in \theta. \quad (5-1)$$

Since $w_{\alpha_P} \alpha \in \Phi_{w^{-1}w_{\alpha_P}}$ and $w^{-1}w_{\alpha_P} \alpha_P = -w^{-1}\alpha_P > 0$, we have by assumption

$$\langle \alpha_P, \alpha^{\vee} \rangle = -\langle \alpha_P, w_{\alpha_P} \alpha^{\vee} \rangle \geq 0. \quad (5-2)$$

From (5-1) and (5-2) it follows that for any $\alpha \in \Phi_{w^{-1}}$ one has

$$\langle D\varpi_P + \rho_M, \alpha^{\vee} \rangle = \sum_{i \in I} m_i \langle \alpha_i, \alpha^{\vee} \rangle \leq 0.$$

This proves that P satisfies property RD. □

Now we present a family of groups with parabolic subgroups satisfying property RD.

Proposition 5.2. *If \mathfrak{g} is a rank 2 hyperbolic Kac–Moody algebra with generalized Cartan matrix $A = \begin{pmatrix} 2 & -b \\ -a & 2 \end{pmatrix}$ where $a, b \geq 2$ and $ab \geq 5$ then every maximal parabolic subgroup of $G_{\mathbb{R}}$ satisfies property RD.*

Proof. First, recall that W is the infinite dihedral group generated by w_1 and w_2 . Without loss of generality, assume that $w^{-1} = (w_2 w_1)^m$ or $w_1 (w_2 w_1)^m$ for some $m \geq 0$. Then each $\alpha \in \Phi_{w^{-1}}$ has the form either

$(w_1w_2)^n\alpha_1$ or $(w_1w_2)^nw_1\alpha_2$ for some $n \geq 0$. As in the proof of [Carbone et al. 2020, Proposition 4.4], define

$$\mu = \frac{\sqrt{ab} + \sqrt{ab-4}}{2} \quad \text{and} \quad h_n = \frac{1}{\mu - \mu^{-1}}(\mu^n - \mu^{-n}).$$

Then we get

$$(w_1w_2)^n\alpha_1 = h_{2n+1}\alpha_1 + \sqrt{a}/\sqrt{b}h_{2n}\alpha_2, \tag{5-3}$$

$$(w_1w_2)^nw_1\alpha_2 = \sqrt{a}/\sqrt{b}h_{2n+2}\alpha_1 + h_{2n+2}\alpha_2. \tag{5-4}$$

Since $w^{-1}\alpha_2 > 0$ and $w^{-1}\alpha_1 < 0$, we need only to show $\langle \alpha_2, \alpha^\vee \rangle < 0$ for $\alpha \in \Phi_{w^{-1}}$ in order to apply Proposition 5.1. Because $\langle \alpha_2, \alpha^\vee \rangle$ and $\langle \alpha, \alpha_2^\vee \rangle$ have the same sign, we consider the latter for convenience. When $\alpha = (w_1w_2)^n\alpha_1$, we obtain from (5-3) and [Carbone et al. 2020, (4.10)]


$$\langle \alpha, \alpha_2^\vee \rangle = \langle h_{2n+1}\alpha_1 + \sqrt{a}/\sqrt{b}h_{2n}\alpha_2, \alpha_2^\vee \rangle = -\sqrt{a}/\sqrt{b}(\sqrt{ab}h_{2n+1} - 2h_{2n}) < 0.$$

Similarly, when $\alpha = (w_1w_2)^nw_1\alpha_2$, we obtain from (5-4) and [Carbone et al. 2020, (4.10)]

$$\langle \alpha, \alpha_2^\vee \rangle = \langle \sqrt{a}/\sqrt{b}h_{2n+2}\alpha_1 + h_{2n+2}\alpha_2, \alpha_2^\vee \rangle = -\sqrt{ab}h_{2n+2} + 2h_{2n+1} < 0.$$

It now follows from Proposition 5.1 that parabolic subgroups satisfy property RD. □

Example 5.3 (nonexample). Consider the following generalized Cartan matrix and its Dynkin diagram:

$$A = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 & -1 \\ 0 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 2 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 2 & -1 \\ 0 & -1 & 0 & 0 & -1 & -1 & 2 \end{pmatrix}$$


Take $\theta = \{1, 2, 3, 4, 5, 6\}$, $w = w_{\alpha_7}w_{\alpha_2}w_{\alpha_1}w_{\alpha_3}w_{\alpha_2}w_{\alpha_7}$ and $\alpha = \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_7$. Then $w \in W^\theta$ and $\alpha \in \Phi_{w^{-1}}$. Since $\langle \varpi_P, \alpha^\vee \rangle = 1$ and $\langle \rho_M, \alpha^\vee \rangle = 0$, we have

$$\langle D\varpi_P + \rho_M, \alpha^\vee \rangle = D.$$

Thus the maximal parabolic subgroup P_θ does not satisfy property RD.

Remark 5.4. Generally, it seems difficult to show that a parabolic subgroup satisfies property RD. However, some computer computations suggest that a large class of Kac–Moody groups have parabolic subgroups with property RD. In Example 5.3, if we take $\theta = \{1, 2, 3, 4, 5, 7\}$, then it appears that P_θ satisfies property RD.

In the rest of this section, we consider the rank 3 hyperbolic Kac–Moody algebra \mathcal{F} studied by Feingold and Frenkel [1983], whose Cartan matrix is

$$A = (a_{ij}) = \begin{pmatrix} 2 & -2 & 0 \\ -2 & 2 & -1 \\ 0 & -1 & 2 \end{pmatrix}.$$

We will see below that the condition of Proposition 5.1 fails for the algebra \mathcal{F} , but there are only finitely many exceptions and we can still show that every maximal parabolic subgroup with finite-dimensional Levi satisfies property RD.

Since A is symmetric, we may identify $\langle \cdot, \cdot \rangle$ with the unique bilinear form $(\cdot | \cdot)$ on \mathfrak{h}^* satisfying $(\alpha_i | \alpha_j) = a_{ij}$, $1 \leq i, j \leq 3$. Define the linear isomorphism of vector spaces $\psi : \mathfrak{h}^* \xrightarrow{\cong} S_2(\mathbb{C})$, the vector space of 2×2 symmetric complex matrices, by

$$\psi(x\alpha_1 + y\alpha_2 + z\alpha_3) = \begin{pmatrix} y-z & y-x \\ y-x & z \end{pmatrix}, \quad (5-5)$$

so that

$$\psi(\alpha_1) = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}, \quad \psi(\alpha_2) = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \quad \psi(\alpha_3) = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$$

and

$$(\lambda | \lambda) = -2 \det \psi(\lambda), \quad \lambda \in \mathfrak{h}^*.$$

For convenience we shall henceforth identify \mathfrak{h}^* with $S_2(\mathbb{C})$ using ψ . Then, for example, the inner product $(\cdot | \cdot)$ is given by

$$\left(\begin{pmatrix} v_1 & v_2 \\ v_2 & v_3 \end{pmatrix} \middle| \begin{pmatrix} \mu_1 & \mu_2 \\ \mu_2 & \mu_3 \end{pmatrix} \right) = -v_3\mu_1 + 2v_2\mu_2 - v_1\mu_3.$$

The Weyl group W of \mathcal{F} is generated by

$$w_1 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad w_2 = \begin{pmatrix} -1 & 1 \\ 0 & 1 \end{pmatrix}, \quad w_3 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad (5-6)$$

and acts on $S_2(\mathbb{C})$ by $g \cdot S = gSg^t$; it is isomorphic to $\mathrm{PGL}_2(\mathbb{Z})$.

The set of real roots is given by

$$\Phi_{\mathrm{re}} = \left\{ \begin{pmatrix} n_1 & n_2 \\ n_2 & n_3 \end{pmatrix} \in S_2(\mathbb{Z}) \mid n_1n_3 - n_2^2 = -1 \right\},$$

and the set of positive real roots is determined to be

$$\Phi_{\mathrm{re}}^+ = \left\{ \begin{pmatrix} n_1 & n_2 \\ n_2 & n_3 \end{pmatrix} \in S_2(\mathbb{Z}) \mid n_1n_3 - n_2^2 = -1, n_1 + n_3 \geq n_2, n_1 + n_3 \geq 0, n_3 \geq 0 \right\}. \quad (5-7)$$

For the rest of this section we write

$$\alpha = \begin{pmatrix} n_1 & n_2 \\ n_2 & n_3 \end{pmatrix} \in \Phi_{\mathrm{re}}^+ \quad \text{and} \quad v = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in W, \quad (5-8)$$

and frequently use the computation

$$v \cdot \alpha = \begin{pmatrix} a^2n_1 + 2abn_2 + b^2n_3 & acn_1 + (ad + bc)n_2 + bdn_3 \\ acn_1 + (ad + bc)n_2 + bdn_3 & c^2n_1 + 2cdn_2 + d^2n_3 \end{pmatrix}. \tag{5-9}$$

In particular,

$$v \cdot \alpha_1 = \begin{pmatrix} -2ab & -bc - ad \\ -bc - ad & -2cd \end{pmatrix}, \tag{5-10}$$

$$v \cdot \alpha_2 = \begin{pmatrix} a^2 + 2ab & ac + bc + ad \\ ac + bc + ad & c^2 + 2cd \end{pmatrix}, \tag{5-11}$$

$$v \cdot \alpha_3 = \begin{pmatrix} b^2 - a^2 & bd - ac \\ bd - ac & d^2 - c^2 \end{pmatrix}. \tag{5-12}$$

Since the elements of Φ_v are precisely the positive real roots flipped by v , a root $\alpha \in \Phi_v$ is constrained by the inequalities

$$c^2n_1 + 2cdn_2 + d^2n_3 \leq 0, \tag{5-13}$$

$$(a^2 + c^2)n_1 + 2(ab + cd)n_2 + (b^2 + d^2)n_3 \leq 0, \tag{5-14}$$

$$(a^2 + c^2)n_1 + 2(ab + cd)n_2 + (b^2 + d^2)n_3 \leq acn_1 + (ad + bc)n_2 + bdn_3, \tag{5-15}$$

in addition to (5-7).

We have two standard maximal parabolic subgroups P_1 and P_2 with finite-dimensional Levi subgroups M_1 and M_2 , which correspond to simple roots α_1 and α_2 respectively. Note that in terms of previous notation, they have semisimple subgroups $L_1 \cong \text{SL}(3)$ and $L_2 \cong \text{SL}(2) \times \text{SL}(2)$ respectively.

We will use the following lemma to prove that the parabolic subgroups P_1 and P_2 satisfy property RD (Proposition 5.6). The proof of this lemma is by direct calculation and will be omitted.

Lemma 5.5. *If $w \in W$, $w \neq \text{id}$ and $w^{-1}\alpha_i > 0$, then $\langle \alpha_i, \alpha^\vee \rangle \leq 0$ for any $\alpha \in \Phi_{w^{-1}}$ unless*

(i) $i = 2$ and α equals

$$\beta_1 = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \quad \text{or} \quad \beta_2 = \begin{pmatrix} 1 & 2 \\ 2 & 3 \end{pmatrix},$$

or

(ii) $i = 3$ and α equals β_1 as above or

$$\beta_3 = \begin{pmatrix} 0 & -1 \\ -1 & 1 \end{pmatrix}.$$

Moreover in these exceptional cases one has $\langle \alpha_i, \alpha^\vee \rangle = 1$.

Proposition 5.6. *The parabolic subgroups P_1 and P_2 both satisfy property RD.*

Proof. Recall that for $j = 1, 2$ and $D > 0$ sufficiently small, one has

$$D\varpi_j + \rho_{M_j} = \sum_i m_i \alpha_i$$

where $m_j < 0$ and $m_i > 0$ for $i \in \theta_j := \{1, 2, 3\} \setminus \{j\}$. Then we need to show that taking smaller D if necessary, one has

$$\langle D\varpi_j + \rho_{M_j}, \alpha^\vee \rangle = \sum_i m_i \langle \alpha_i, \alpha^\vee \rangle < 0$$

for any $w \in W^{\theta_j}$ and $\alpha \in \Phi_{w^{-1}}$. Note that $\langle \alpha_i, \alpha^\vee \rangle$, $i = 1, 2, 3$, cannot be all zero. We shall consider P_1 and P_2 separately.

(1) Assume that $j = 1$. Then $w^{-1}\alpha_1 < 0$ and $w^{-1}\alpha_i > 0$ for $i = 2, 3$. By Lemma 5.5 and (5-2), one always has $\langle \alpha_1, \alpha^\vee \rangle \geq 0$ for $\alpha \in \Phi_{w^{-1}}$, and it suffices to consider the exceptional cases when $\alpha = \beta_i$ for some $i = 1, 2, 3$.

Since $\beta_1 = \alpha_2 + \alpha_3$, we have $w^{-1}\beta_1 > 0$, hence $\alpha = \beta_1$ cannot happen. Since $\langle \alpha_3, \beta_2^\vee \rangle = 2$, it follows from Lemma 5.5 that $\alpha = \beta_2$ cannot happen either. Finally if $\alpha = \beta_3$, then $\langle \rho_{M_1}, \beta_3^\vee \rangle = -2 < 0$, hence $\langle D\varpi_1 + \rho_{M_1}, \beta_3^\vee \rangle = D - 2 < 0$ for $D < 2$.

(2) Assume that $j = 2$. Then $w^{-1}\alpha_2 < 0$ and $w^{-1}\alpha_i > 0$ for $i = 1, 3$. By Lemma 5.5 and (5-2), one has $\langle \alpha_2, \alpha^\vee \rangle \geq 0$ for $\alpha \in \Phi_{w^{-1}}$ unless $\alpha = w_2\beta_1$ or $w_2\beta_2$. However, we find that $w_2\beta_1 = \alpha_3$ and $\langle \alpha_3, w_2\beta_2^\vee \rangle = 3$, hence $\alpha \neq w_2\beta_1, w_2\beta_2$, thanks to $w^{-1}\alpha < 0$ and Lemma 5.5 respectively. Thus we have proved that $\langle \alpha_2, \alpha^\vee \rangle \geq 0$ always holds.

By Lemma 5.5 again, it suffices to consider the exceptional cases $\alpha = \beta_1$ or β_3 . For $\alpha = \beta_1$ we have $\langle \rho_{M_2}, \beta_1^\vee \rangle = -1/2 < 0$, hence $\langle D\varpi_2 + \rho_{M_2}, \beta_1^\vee \rangle = D - 1/2 < 0$ for $D < 1/2$. The case $\alpha = \beta_3$ cannot happen, because $\langle \alpha_1, \beta_3^\vee \rangle = 2$ which violates Lemma 5.5. \square

Acknowledgments

We are grateful to Stephen D. Miller for his useful comments, and would like to thank the anonymous referees for helpful comments.

References

- [Borel 2007] A. Borel, “Automorphic forms on reductive groups”, pp. 7–39 in *Automorphic forms and applications*, edited by P. Sarnak and F. Shahidi, IAS/Park City Math. Ser. **12**, Amer. Math. Soc., Providence, RI, 2007. MR Zbl
- [Braverman and Kazhdan 2013] A. Braverman and D. Kazhdan, “Representations of affine Kac–Moody groups over local and global fields: a survey of some recent results”, pp. 91–117 in *European Congress of Mathematics*, edited by R. Latafa et al., Eur. Math. Soc., Zürich, 2013. MR Zbl
- [Bump et al. 1996] D. Bump, S. Friedberg, and J. Hoffstein, “On some applications of automorphic forms to number theory”, *Bull. Amer. Math. Soc. (N.S.)* **33**:2 (1996), 157–175. MR Zbl
- [Carbone and Garland 2003] L. Carbone and H. Garland, “Existence of lattices in Kac–Moody groups over finite fields”, *Commun. Contemp. Math.* **5**:5 (2003), 813–867. MR Zbl
- [Carbone et al. 2017] L. Carbone, K.-H. Lee, and D. Liu, “Eisenstein series on rank 2 hyperbolic Kac–Moody groups”, *Math. Ann.* **367**:3-4 (2017), 1173–1197. MR Zbl
- [Carbone et al. 2020] L. Carbone, H. Garland, K.-H. Lee, D. Liu, and S. D. Miller, “On the convergence of Kac–Moody Eisenstein series”, 2020. arXiv 2005.13636
- [De Medts et al. 2009] T. De Medts, R. Gramlich, and M. Horn, “Iwasawa decompositions of split Kac–Moody groups”, *J. Lie Theory* **19**:2 (2009), 311–337. MR Zbl

- [Feingold and Frenkel 1983] A. J. Feingold and I. B. Frenkel, “A hyperbolic Kac–Moody algebra and the theory of Siegel modular forms of genus 2”, *Math. Ann.* **263**:1 (1983), 87–144. MR Zbl
- [Garland 1999] H. Garland, “Eisenstein series on arithmetic quotients of loop groups”, *Math. Res. Lett.* **6**:5-6 (1999), 723–733. MR
- [Garland 2004] H. Garland, “Certain Eisenstein series on loop groups: convergence and the constant term”, pp. 275–319 in *Algebraic groups and arithmetic*, edited by S. G. Dani and G. Prasad, Tata Inst. Fund. Res., Mumbai, 2004. MR
- [Garland 2006] H. Garland, “Absolute convergence of Eisenstein series on loop groups”, *Duke Math. J.* **135**:2 (2006), 203–260. MR
- [Garland 2007a] H. Garland, “Eisenstein series on loop groups: Maass–Selberg relations, I”, pp. 275–300 in *Algebraic groups and homogeneous spaces*, edited by V. B. Mehta, Tata Inst. Fund. Res. Stud. Math. **19**, Mumbai, 2007. MR
- [Garland 2007b] H. Garland, “Eisenstein series on loop groups: Maass–Selberg relations, II”, *Amer. J. Math.* **129**:3 (2007), 723–784. MR
- [Garland 2007c] H. Garland, “Eisenstein series on loop groups: Maass–Selberg relations, III”, *Amer. J. Math.* **129**:5 (2007), 1277–1353. MR Zbl
- [Garland 2007d] H. Garland, “Eisenstein series on loop groups: Maass–Selberg relations, IV”, pp. 115–158 in *Lie algebras, vertex operator algebras and their applications*, edited by Y.-Z. Huang and K. C. Misra, Contemp. Math. **442**, Amer. Math. Soc., Providence, RI, 2007. MR Zbl
- [Garland 2011] H. Garland, “On extending the Langlands–Shahidi method to arithmetic quotients of loop groups”, pp. 151–167 in *Representation theory and mathematical physics*, edited by B. L. Jeffrey Adams and S. Sahi, Contemp. Math. **557**, Amer. Math. Soc., Providence, RI, 2011. MR
- [Garland et al. 2017] H. Garland, S. D. Miller, and M. M. Patnaik, “Entirety of cuspidal Eisenstein series on loop groups”, *Amer. J. Math.* **139**:2 (2017), 461–512. MR
- [Godement 1964] R. Godement, “Domaines fondamentaux des groupes arithmétiques”, exposé 257 in *Séminaire Bourbaki, 1962/1963*, W. A. Benjamin, Amsterdam, 1964. MR Zbl
- [Green et al. 2010] M. B. Green, S. D. Miller, J. G. Russo, and P. Vanhove, “Eisenstein series for higher-rank groups and string theory amplitudes”, *Commun. Number Theory Phys.* **4**:3 (2010), 551–596. MR Zbl
- [Kac 1990] V. G. Kac, *Infinite-dimensional Lie algebras*, 3rd ed., Cambridge University Press, 1990. MR Zbl
- [Kac and Peterson 1985] V. G. Kac and D. H. Peterson, “Defining relations of certain infinite-dimensional groups”, pp. 165–208 in *The mathematical heritage of Élie Cartan* (Lyon, 1984), Astérisque (numéro hors série), Société Mathématique de France, Paris, 1985. MR Zbl
- [Kapranov 2000] M. Kapranov, “The elliptic curve in the S-duality theory and Eisenstein series for Kac–Moody groups”, 2000.
- [Kim 2003] H. H. Kim, “Functoriality for the exterior square of GL_4 and the symmetric fourth of GL_2 ”, *J. Amer. Math. Soc.* **16**:1 (2003), 139–183. MR
- [Kim and Shahidi 2002] H. H. Kim and F. Shahidi, “Functorial products for $GL_2 \times GL_3$ and the symmetric cube for GL_2 ”, *Ann. of Math. (2)* **155**:3 (2002), 837–893. MR Zbl
- [Kumar 2002] S. Kumar, *Kac–Moody groups, their flag varieties and representation theory*, Progress in Mathematics **204**, Birkhäuser, Boston, 2002. MR Zbl
- [Langlands 1976] R. P. Langlands, *On the functional equations satisfied by Eisenstein series*, Lecture Notes in Mathematics **544**, Springer, 1976. MR
- [Lee and Lombardo 2014] K.-H. Lee and P. Lombardo, “Eisenstein series on affine Kac–Moody groups over function fields”, *Trans. Amer. Math. Soc.* **366**:4 (2014), 2121–2165. MR Zbl
- [Liu 2015] D. Liu, “Eisenstein series on loop groups”, *Trans. Amer. Math. Soc.* **367**:3 (2015), 2079–2135. MR Zbl
- [Looijenga 1980] E. Looijenga, “Invariant theory for generalized root systems”, *Invent. Math.* **61**:1 (1980), 1–32. MR Zbl
- [Miller and Schmid 2012] S. D. Miller and W. Schmid, “On the rapid decay of cuspidal automorphic forms”, *Adv. Math.* **231**:2 (2012), 940–964. MR Zbl
- [Mœglin and Waldspurger 1995] C. Mœglin and J.-L. Waldspurger, *Spectral decomposition and Eisenstein series*, Cambridge Tracts in Mathematics **113**, Cambridge University Press, 1995. MR

[Patnaik 2008] M. M. Patnaik, *Geometry of loop Eisenstein series*, ProQuest LLC, Ann Arbor, MI, 2008. MR

[Shahidi 2005] F. Shahidi, “Infinite dimensional groups and automorphic L -functions”, *Pure Appl. Math. Q.* **1:3** (2005), 683–699. MR Zbl

Communicated by Peter Sarnak

Received 2020-08-26 Revised 2021-05-08 Accepted 2021-08-25

carbonel@math.rutgers.edu

*Department of Mathematics, Rutgers, The State University of New Jersey,
Piscataway, NJ, United States*

khlee@math.uconn.edu

*Department of Mathematics, University of Connecticut, Storrs, CT,
United States*

maliu@zju.edu.cn

School of Mathematical Sciences, Zhejiang University, Hangzhou, China

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR
Antoine Chambert-Loir
Université Paris-Diderot
France

EDITORIAL BOARD CHAIR
David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Philippe Michel	École Polytechnique Fédérale de Lausanne
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Frank Calegari	University of Chicago, USA	Raman Parimala	Emory University, USA
J.-L. Colliot-Thélène	CNRS, Université Paris-Saclay, France	Irena Peeva	Cornell University, USA
Brian D. Conrad	Stanford University, USA	Jonathan Pila	University of Oxford, UK
Samit Dasgupta	Duke University, USA	Anand Pillay	University of Notre Dame, USA
Hélène Esnault	Freie Universität Berlin, Germany	Bjorn Poonen	Massachusetts Institute of Technology, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Victor Reiner	University of Minnesota, USA
Sergey Fomin	University of Michigan, USA	Peter Sarnak	Princeton University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Wee Teck Gan	National University of Singapore	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	Shunsuke Takagi	University of Tokyo, Japan
Ben J. Green	University of Oxford, UK	Pham Huu Tiep	Rutgers University, USA
Christopher Hacon	University of Utah, USA	Ravi Vakil	Stanford University, USA
Roger Heath-Brown	Oxford University, UK	Akshay Venkatesh	Institute for Advanced Study, USA
János Kollár	Princeton University, USA	Melanie Matchett Wood	Harvard University, USA
Michael J. Larsen	Indiana University Bloomington, USA	Shou-Wu Zhang	Princeton University, USA

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2022 is US \$450/year for the electronic version, and \$655/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

ANT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2022 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 16 No. 5 2022

Artin's conjecture for Drinfeld modules WENTANG KUO and DAVID TWEEDLE	1025
The G -stable rank for tensors and the cap set problem HARM DERKSEN	1071
Entirety of certain cuspidal Eisenstein series on Kac–Moody groups LISA CARBONE, KYU-HWAN LEE and DONGWEN LIU	1099
Resolution of ideals associated to subspace arrangements ALDO CONCA and MANOLIS C. TSAKIRIS	1121
On the geometry of geometric rank RUNSHI GENG and JOSEPH M. LANDSBERG	1141
The average size of the 2-Selmer group of a family of non-hyperelliptic curves of genus 3 JEF LAGA	1161
Multiplicities and Betti numbers in local algebra via lim Ulrich points SRIKANTH B. IYENGAR, LINQUAN MA and MARK E. WALKER	1213
The mean values of cubic L -functions over function fields CHANTAL DAVID, ALEXANDRA FLOREA and MATILDE LALÍN	1259